



Kelvin Open Science Publishers
Connect with Research Community

Research Article

Volume 1 / Issue 1

KOS Journal of Environmental and Earth Science

<https://kelvinpublishers.com/journals/environmental-and-earth-science.php>

Sustainable Use of Nature Resources and Agriculture High-Quality Development

Zhongsheng Guo^{1,2,*}

¹Institute of Water and Soil Conservation, Northwest A&F University, China

²Department of Water Resources, Institute of Water and Soil Conservation, Chinese Academy of Sciences, China

*Corresponding author: Zhongsheng Guo, Institute of Water and Soil Conservation, Northwest A&F University, China

Received: March 28, 2025; **Accepted:** April 03, 2025; **Published:** April 05, 2025

Citation: Zhongsheng G. (2025) Sustainable Use of Nature Resources and Agriculture High-Quality Development. *J Envi Earth Sci*. 1(1): 1-4.

Copyright: © 2025 Zhongsheng G., This is an open-access article published in *J Envi Earth Sc* and distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Abstract

As economy develops and population increase, there is an increasing demand for plant goods such as timber, food, medicine and so on and ecological service such as clean air, fresh water and environment. Because plant goods and service producing by original forest cannot meet the increasing need of people, most of the original forest has gradually become fruit, farmland, plantation and grass. A lot of exotic plant was introduced to produce special production and service in the plant restoration. As introduced plant grow, soil and vegetation degradation and crop failure will happen because plant overload and overuse nature resources or plant waste resources because plant underuse nature resources because underload. In order to solve these problems, according to my years of research, the results have showed that there is a resource use limit by plants and vegetation carrying capacity. Resources use limit by plants includes space resources use limit by plants in soil water and nutrient rich regions, soil water resources use limit by plants in water-limited regions and soil nutrient resources use limit by plants in nutrient-limited regions. Vegetation carrying capacity includes space vegetation carrying capacity, soil water vegetation carrying capacity and soil nutrient vegetation carrying capacity in the process of plant growth. When the available amount of nature resources reduced to resources use limit by plants, such as soil water resources use limit by plants, which is the soil water resources in the range of the maximum infiltration depth is equal to soil water resources use limit by plants, the plant resources relation enters the key period of plant resources relation regulation. The ending time of the key period of plant resources relation regulation is the ineffective time of plant resources relation regulation. If the existing plant density is more than vegetation carrying capacity in the key period of plant resources relation regulation, the plant resources relation must be regulated on the vegetation carrying capacity to get maximal yield and beneficial effect and realize sustainable use of nature resources, high quality sustainable management of forest vegetation and agriculture high quality production.

2. Keywords

Key period of plant resources relation regulation, Plant grow, Resources use limit by plants, Vegetation carrying capacity, High quality sustainable management of forest vegetation,

Agriculture high quality production

3. Introduction

The high-quality development of agriculture is to take some measures and methods to make the land produce the

maximum output and services to meet people's yearning for a better life and the needs of agricultural production services.

As economy develops and population increases, there is an increasing demand for the quantity and variety of plant production and service, but the plant production, especially food, fruit, fibre and wood and so on produced by original forest ecosystem cannot meet the need of demand for the quantity and variety of timber, food, medicine and so on and ecological service, so 80% of the original forest in the loess plateau [1] and the world has become farmland, man-made forest and grass [2,3] and a lot of exotic plant was introduced to produce more production and service. Because exotic plant changes the plant resources relation and the ability of self-regulation to adapt to climate change is very low, vegetation decline, crop failure, serious soil erosion, greenhouse gas emissions and other ecological and environmental problems happens [4].

In order to solve the problem of soil and vegetation degradation or crop failure or waste of resources and achieve high-quality development and meet the increasing need of people for plant goods and service. Years of research has shown that there is a limit of plant utilization of soil water resources and the carrying capacity of soil water to vegetation the method of high quality and sustainable management of forest vegetation in water shortage area in the process of vegetation restoration [2,3,5]. The main contents are as follows:

3.1. The utilization limit of soil water resources


Along with plant growth, plant height, diameter and canopy volume increase, and roots develop deep and soil water resources in the root zone reduced but available soil water resources are declined. In order to obtain maximum yield and beneficial result and achieve high quality and sustainable development, even the root distribution depth is more than the maximum infiltration depth, see Figure 1, plant roots do not absorb water indefinitely due to lack of water such as in semiarid loess hilly region (Guyuan, China) the precipitation changes with year and month, see Figure 2 [1-3,6]. There should be a control limit for the utilization of soil water resources by plants in water-scarce areas, that is, the soil water resources use limit by plants [7,8]. It is the soil water storage when the soil water content is equal to the wilting coefficient in the range of the maximum infiltration depth. The wilting coefficient at different soil depth in the maximum infiltration depth can be determined by centrifugal machine or press film. The undisturbed soil at different soil depth in a soil profile was taken by a cutting ring with a 5 cm high and 5 cm diameter. The maximum infiltration depth can be estimated by the two curves method [9,10]. The wilting coefficient is expressed by the wilting coefficient of indicating plants in a plant community. The indicator plants of natural vegetation are the constructive species, and the artificial vegetation is the target species. When the soil water resources decrease to the utilization limit of soil water resources, the plant-water relationship enters the starting time of critical period of plant-water relationship regulation [11,12]. At this time, the regulation of plant-water relationship should be considered to be regulated. The initial planting density influences the soil water resource use limit by plants [13].

Figure 1: The *Robinia pseudoacacia* L. root distribution depth and maximum infiltration depth (290 cm) relationship

in semiarid loess hilly region (Guyuan, Ningxia, China).



Figure 2: Change of precipitation with month (photo above) and year (photo below) in semiarid loess hilly region (Guyuan, China).



3.2. Vegetation carrying capacity

Vegetation carrying capacity is the ability of land resources to support vegetation, which is limited. The vegetation carrying capacity in water shortage area is the soil water vegetation carrying capacity, which is the ability of soil water resources to support vegetation and is limited. The soil water vegetation carrying capacity refers to the maximum quantity per unit area of soil water resources that can maintain healthy growth of indicator plants at a given period and site condition [2,3,9,14,15] and can be estimated by soil water-plant density model [2,3,9], expressed by the number (absolute index) or density (relative index) of indicator plant population in the plant community. Soil water vegetation carrying capacity (SWVCC), changing with plant species (vegetation type), time or position, time climate change, especially the critical period of plant water relationship regulation, which is the theoretical basis for determining indicators and criteria for forest resources use degree and high quality and sustainable management of forest vegetation [2,3,9].

3.3. Critical period of plant-resources relationship regulation

Generally, along with plant growth, the available solar resources amount is limit in a given canopy (cube) such as leaf amount in a given canopy or soil water resources in the maximum infiltration depth will also decrease, even though the soil water resources will increase after the rain. The plant water relationship enters the critical period of plant water relationship regulation. If the present plant density is more than soil water vegetation carrying capacity in the critical period of plant water relationship regulation, plant water relationship regulation has to be regulated. Plant-water relationship can be regulated plant-water relationship can be regulated by different methods such as cultivating and selecting better plant species and varieties [5], soil preparations such as thin film cover and ridge tillage in the semiarid loess hilly region (Guyuan, China), see Figure 3 and pruning some leave and trig and so on, but the most important method is to regulate the plant-water relationship by reducing plant density in the critical period of plant-water relationship regulation.

Figure 3: Photo of corn thin film cover and ridge tillage in semiarid loess hilly region on 21 May, 2021.



Plant-resources relationship in the growing season can be divided into different stages according to resources use limit by plants such as soil water resources use limit by plants in the water-limited regions. If the available resources in the canopy are more resources use limit by plants or soil water resources in the range of the maximum infiltration depth is more than soil water resources use limit by plants, showing resources such as soil water resources enough and plant grow healthily. If not, more attention should be paid to the plant-water relationship regulation.

The critical period of plant-resources relationship regulation refers to the time that the resources can maintain the plant growth healthy when the resources descend to the utilization limit of resources by plants such as soil water resources within the range of the maximum infiltration depth, it not only affects plant growth condition but also determines the maximum yield and benefit in the growing season in the water-limited regions. The critical period of plant water relationship regulation is the period from the starting time soil water resources reduced to the limit of soil water resources utilization by plants to the ending time plant water relationship regulation is failure. The ending time of critical period of plant water relationship regulation can be estimate by the thinning plant method. Sowing seeds in a plot with a maximum plant density and ensure the maximum experimental plant density equals to or more than soil water carrying capacity for vegetation.

3.4. High quality development of agriculture

Firstly, the better plant species and varieties have to be selected according to site condition and cultivation goal [5] and then suitable initially planting density has to be taken. The initially planting density is more than vegetation carrying capacity such as soil water vegetation carrying capacity in water-limited regions. If the existing plant density is more than soil water carrying capacity for vegetation in the critical period of plant-water relationship regulation, showing plant overuse soil water and plant density exceeds soil water carrying capacity, which will cause soil degradation, vegetation decline, fruit and crop failure if plant-water relationship should not be regulated. At this movement, we should regulate the plant-water relationship by reducing the plant density. The reducing amount of plant density is equal to the different between the existing plant density and soil water carrying capacity for vegetation. When the soil water resources decrease to the utilization limit of soil water resources by plants in the range of maximum infiltration depth, plant-water relationship enters the critical period of plant-water relationship regulation. If the existing plant density is more than soil water carrying capacity for vegetation in the critical period of plant-water relationship regulation, plant-water relationship should be regulated by estimating the soil water carrying capacity for vegetation in the critical period of plant-water relationship regulation, and the maximum yield and benefit of non-commercial forest and grass can be obtained by determining and controlling the plant-water relationship according to the soilwater carrying capacity for vegetation in the critical period of plant-water relationship regulation. The high quality and sustainable development can be achieved in water-limited regions.

4. Conclusion

The most important work of forest restoration is the plant-resources relationship regulation in the process of vegetation according to the cultivated goal, especially plant-resources relationship regulation in the critical period of plant-resources relationship regulation such because it decides the maximum yield and benefit of a population or plant community. The foundation of plant-resources relationship regulation is resources use limit by plants. When resources amount reduced to utilization limit of resources by plants, such as the soil water resources within the range of the maximum infiltration depth descend to the utilization limit of soil water resources, plant-resources relationship enters the critical period of plant-resources relationship regulation in which the resources severe influence the plant growth and decides the maximum yield and ecological, economic and social benefits. At this time, the plant-resources relationship should be regulated to obtain the maximum yield and benefit of non-commercial forest and grass and carry out agriculture high quality development.

5. Acknowledgement

This research was supported by the National Science Fund of China (Project No's 42077079, 41071193 and 41271539) and study on high quality sustainable development of soil and moisture conservation (A2180021002).

6. References

1. Guo Z, Shao M (2013) Impact of afforestation density on soil and water conservation of the semiarid loess plateau, China. *J Soil Water Conserv.* 68(5): 401-410.
2. Guo ZS (2021) Soil water carrying capacity for vegetation. *Land Degradation Development.* 32(14): 3801-3811.

3. Guo Z (2021) Soil hydrology process and rational use of soil water resources in desert regions. *Water*. 13(17): 2377.
4. Chen H, Shao M, Li Y (2008) Soil desiccation in the loess plateau of China. *Geoderma*. 143: 91-100.
5. Guo ZS, Wu QX, Si LM (1990) A study on selecting trees and grasses species of woodland for fuel, fodder and manure using fuzzy mathematics method in the loess hilly area. *Journal of Biomathematics*. 5(3): 69-76.
6. Guo Z (2019) Rice carrying capacity and sustainable produce of rice in resources-limited regions. *Int J Agric Sc Food Technol*. 5(1): 54-57.
7. Guo Z, Shao M (2010) Effect of artificial *Caragana korshinskii* forest on soil water in the semiarid area of loess hilly region. *Chin Forest Sci*. 46: 1-8.
8. Guo Z (2010) Soil water resource use limit in semi-arid loess hilly area. *Ying Yong Sheng Tai Xue Bao*. 21: 3029-3035.
9. Guo Z (2014) Theory and practice on soil water carrying capacity for vegetation. *Chin Scientific Press*. 45-100.
10. Guo Z (2020) Estimating method of maximum infiltration depth and soil water supply. *Sci Rep*. 10(1): 9726.
11. Guo Z (2009) Limit of vegetation rehabilitation for soil and water conservation in semi-arid region of loess plateau: A case study of artificial *Caragana korshinskii* stand. *Science of Soil and Water Conservation*. 7: 49-54.
12. Guo Z, Li Y (2009) Initiation stage to regulate the caragana growth and soil water in the semiarid area of loess hilly region, China. *Acta Ecologica Sinica*. 29: 5721-5729.
13. Guo Z, Zhang W (2016) Impact of initial planting density on soil water resource use limit by plants. *Geoinfor Geostat: An Overview*. 4: 1.
14. Guo Z, Shao M, Zhangy, Wu Q (2002) An layer-dividing approach to the soil water in forest land, the proceedings of soil physics and ecological environmental construction. *Xiaan, Shanxi Science and Technology Press*. 74-79.
15. Guo Z, Shao M (2003) Vegetation carrying capacity of soil water and soil desiccation in artificial forestry and grassland in the semiarid regions of loess plateau. *Chin J Ecol*. 23: 1640-1647.